arXiv:math-ph/0608021v2 13 Dec 2007

Journal of Mathematical Physics

Polynomial Poisson Algebras for Classical Superintegrable Systems with a Third Order Integral of Motion

Ian Marquette
Département de physique et Centre de recherche mathématiques, Université de Montréal,
C.P.6128, Succursale Centre-Ville, Montréal, Québec H3C 3J7, Canada
ian.marquette@umontreal.ca
514-343-6111 (4072)

Pavel Winternitz
Département de mathématiques et de statistique et Centre de recherche mathématiques,Université de Montréal,
C.P.6128, Succursale Centre-Ville, Montréal, Québec H3C 3J7, Canada
wintern@CRM.UMontreal.CA
514-343-7271

We present polynomial Poisson algebras for the 8 classical potentials in two-dimensional Euclidian space that separate in cartesian coordinates and allow a third order integral of motion. Some of the classical superintegrable potentials do not coincide with quantum ones, but are their singular limits. We present the trajectories for all these classical potentials. We find that all bounded trajectories are periodic.

1 Introduction

The purpose of this article is to study the properties of a certain class of superintegrable systems in a real two-dimensional Euclidian space E_2. These are classical hamiltonian systems that allow three functionally independent integrals of motion. One of them is the Hamiltonian H. The other two are second and third order polynomials in the momenta, respectively. Moreover, in most of the article we shall restrict to the case when the potential allows separation of variables in Cartesian coordinates, so that we have

$$H = \frac{1}{2}(P_1^2 + P_2^2) + f(x) + g(y) \tag{1.1}$$

$$A = \frac{1}{2}(P_1^2 - P_2^2) + f(x) - g(y) \tag{1.2}$$

and X_3 will have the general form

$$X_3 = \sum_{i+j+k=3} a_{ijk} L_3^i P_1^j P_2^k + k_1(x,y)P_1 + k_2(x,y)P_2, \tag{1.3}$$

where a_{ijk} are constant and L_3 is the angular momentum

$$L_3 = xP_2 - yP_1. \tag{1.4}$$

Superintegrable systems are Hamiltonian systems with more integrals of motion than degrees of freedom. In dimension n=2 the maximal possible number of functionally independent integrals is 2n-1=3 and that is the case we are considering.

A large body of literature exists on superintegrable systems, inspired by the superintegrability of the harmonic oscillator[1,2] and Kepler-Coulomb potentials[3,4]. A systematic search for superintegrable systems in E_2 and E_3 was conducted in the 1960ties[5,6,7]. Like much of the later work on superintegrable systems it was restricted to the case of second order integrals of motion[8,..,13]. This case turned out to have an intimate connection with the separation of variables in the Hamilton-Jacobi and Schrödinger equations.

An interesting discovery was that in general the quadratic integrals of

motion do not generate finite dimensional Lie algebras, but more complicated algebraic structures, namely quadratic algebras[14−18].
Much less is known about integrable and superintegrable systems with third and higher order integrals of motion. In 1935 Drach found 10 different potentials in a complex Euclidian plane allowing a third order integral of motion in classical mechanics[19,20]. In 1984 Hietarinta[21] showed that in the case of integrals that are third, or higher order polynomials in momenta, quantum and classically integrable potentials may not coincide.
A systematic search for superintegrable systems involving at least one third order integral was started in 2002[22]. It was shown that in the case of one first order and one third order integral the only classical superintegrable systems were known ones (like $V = \alpha r^2$ or $V = \alpha r^{-1}$) for which the third order integral was the product of a first and second order one. However, in the quantum case a new superintegrable potential of this type exists, namely $V_1(x, y) = (\hbar\omega)^2 k^2 sn^2(\omega x, k)$ where $sn(\omega x, k)$ is a Jacobi elliptic function. The classical limit ($\hbar \to 0$) is free motion ![22]
All classical and all quantum potentials allowing a second order integral of the form (1.2) and a third order integral (of any form) were found in ref.[23]. The classical ones are all expressed in terms of elementary functions. The quantum ones are quite different and some of them involve elliptic functions and Painlevé transcendents[24]. We leave the quantum case for a future study and concentrate here on the classical superintegrable Hamiltonian systems of the form (1.1),...,(1.3)

In Section 2 we consider a general superintegrable Hamiltonian system in a two-dimensional space (not necessarily Euclidian) with a scalar potential. It allows one quadratic and one cubic integral of motion. We construct the most general cubic Poisson algebra generated by these integrals. We express the Casimir operator K of this algebra in terms of the Hamiltonian H. The general formalism of Section 2 is applied in Section 3 to the 8 existing superintegrable systems in E_2 allowing separation of variables in Cartesian coordinates and having an additional third order integral of motion. Solutions of the equation of motion and trajectories in these 8 systems are discussed in Section 4. Section 5 is devoted to conclusions.

2 Cubic Poisson algebras

We start out with a more general Hamiltonian system than (1.1)...(1.3). We will not assume that we have a natural Hamiltonian in E_2, but that we have a superintegrable system with a quadratic Hamiltonian and one second order and one third order integral of motion. We put

$$H = a(q_1,q_2)P_1^2 + 2b(q_1,q_2)P_1P_2 + c(q_1,q_2)P_2^2 + V(q_1,q_2)$$

$$\begin{aligned} A = A(q_1,q_2,P_1,P_2) = d(q_1,q_2)P_1^2 + 2e(q_1,q_2)P_1P_2 + \\ f(q_1,q_2)P_2^2 + g(q_1,q_2)P_1 + h(q_1,q_2)P_2 + Q(q_1,q_2) \end{aligned} \tag{2.1}$$

$$B = B(q_1,q_2,P_1,P_2) = u(q_1,q_2)P_1^3 + 3v(q_1,q_2)P_1^2P_2 + 3w(q_1,q_2)P_1P_2^2$$

$$+x(q_1,q_2)P_2^3 + j(q_1,q_2)P_1^2 + 2k(q_1,q_2)P_1P_2 + l(q_1,q_2)P_2^2 + m(q_1,q_2)P_1 + n(q_1,q_2)P_2 + s(q_1,q_2)$$

where P_i and q_i are canonical momenta and coordinates and a,..,s are functions to be determined. Since A and B are integrals of motion we have.

$$\{H,A\} = \{H,B\} = 0 \tag{2.2}$$

where $\{,\}$ is the Poisson bracket

Since B is cubic in the momenta, we cannot expect $\{H, A$ and B$\}$ to generate a quadratic algebra, like those obtained in[14,...,18]. We will however try to close the algebra at the lowest order possible, namely 3. We put

$$\begin{aligned} \{A,B\} &= C \\ \{A,C\} &= \alpha A^2 + 2\beta AB + \gamma A + \delta B + \epsilon \\ \{B,C\} &= \mu A^3 + \nu A^2 + \rho B^2 + 2\sigma AB + \xi A + \eta B + \zeta \end{aligned} \tag{2.3}$$

Since we have $\{C,\{A,B\}\}$=0 the Jacobi identity reduces to

$$\{A,\{B,C\}\} = \{B,\{A,C\}\} \tag{2.4}$$

The Jacobi identity implies

$\rho = -\beta, \sigma = -\alpha, \eta = -\gamma$ and we obtain the cubic algebra.

$$\begin{aligned} \{A, B\} &= C \\ \{A, C\} &= \alpha A^2 + 2\beta AB + \gamma A + \delta B + \epsilon \\ \{B, C\} &= \mu A^3 + \nu A^2 - \beta B^2 - 2\alpha AB + \xi A - \gamma B + \zeta. \end{aligned} \tag{2.5}$$

The coefficients α , β and μ are constants, but the other ones can be polynomials in the Hamiltonian H. The degrees of these polynomials are dictated by the fact that H and A are second order polynomials in the momenta and B is a third order one. Hence C can be a fourth order polynomial. We have

$$\begin{gathered} \alpha = \alpha_0, \beta = \beta_0, \mu = \mu_0 \\ \gamma = \gamma_0 + \gamma_1 H, \delta = \delta_0 + \delta_1 H, \epsilon = \epsilon_0 + \epsilon_1 H + \epsilon_2 H^2 \\ \nu = \nu_0 + \nu_1 H, \xi = \xi_0 + \xi_1 H + \xi_2 H^2 \\ \zeta = \zeta_0 + \zeta_1 H + \zeta_2 H^2 + \zeta_3 H^3, \end{gathered} \tag{2.6}$$

where α_0,...,ζ_3 are constants. A Casimir operator K of a polynomial algebra is defined as an operator Poisson commuting with all elements of the algebra. For the algebra (2.5) this means

$$\{K, A\} = \{K, B\} = \{K, C\} = 0 \tag{2.7}$$

and this implies

$$K = C^2 - 2\alpha A^2 B - 2\beta AB^2 - 2\gamma AB - \delta B^2 - 2\epsilon B + \frac{1}{2}\mu A^4 + \frac{2}{3}\nu A^3 + \xi A^2 + 2\zeta A. \tag{2.8}$$

Thus K is a polynomial of order 8 in the momenta. since the Hamiltonian H also satisfies relations (2.7) we can expect K to be a polynomial in H and we write

$$K = k_0 + k_1 H + k_2 H^2 + k_3 H^3 + k_4 H^4 \tag{2.9}$$

where $k_0, .., k_4$ are constants. Note that eq. (2.9) together with (2.8) represents a polynomial relation between the integrals H, A, B and C in agreement with the fact that only 3 of them can be functionally independent.

3 Cubic Poisson algebras for classical superintegrable systems

Let us now apply the formalism of Section 2 to all classical superintegrable potentials that separate in Cartesian coordinates and allow a third order integral of motion[22,23]. They have the form (1.1),...,(1.3) , so we have $a(q_1, q_2) = c(q_1, q_2) = 1$, $b(q_1, q_2) = 0$ in (2.1). Eight such systems exist, four of them are well-known, four first presented in[23].

Case 1. The isotropic harmonic oscillator.
We have

$$H = \frac{P_1^2}{2} + \frac{P_1^2}{2} + \frac{\omega^2}{2}(x^2 + y^2) \tag{3.1}$$

The formalism of Section 2 applies, but we cannot expect it to provide anything new. indeed, in this case we have a well known u(2) algebra of first and second order integrals of motion :

$$\begin{gathered} L_3 = xP_2 - yP_1, X_1 = P_1^2 + \omega^2 x^2, X_2 = P_2^2 + \omega^2 y^2 \\ X_3 = P_1 P_2 + \omega^2 xy \end{gathered} \tag{3.2}$$

we can chose

$$\begin{gathered} A = P_1^2 - P_2^2 + \omega^2(x^2 - y^2) = X_1 - X_2 \\ B = L_3 X_2 = xP_2^3 - yP_1P_2 - \omega^2 y^3 P_1 + \omega^2 xy^2 P_2 \end{gathered} \tag{3.3}$$

we obtain

$$\{A, B\} = C = -4X_2X_3$$

$$\{A,C\} = -16\omega^2 B \tag{3.4}$$
$$\{B,C\} = 2A^3 - 6HA^2 + 8H^3$$

Thus, relations (2.5) are satisfied, but we remain in the enveloping algebra of u(2). The Casimir operator of the cubic algebra in this case is

$$K = C^2 + 16\omega^2 B^2 + 2A^4 - 6HA^3 + 8H^3 A = 16H^4 \tag{3.5}$$

However, in this case H is simply the central element of the u(2) algebra (3.2) and it generates the center of the enveloping algebra of u(2)(a hence also of the cubic algebra).

Case 2. A quadratically superintegrable Hamiltonian[5,6]

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + \frac{\omega^2}{2}(x^2+y^2) + \frac{b}{x^2} + \frac{c}{y^2} \tag{3.6}$$

The Hamiltonian allows two second order integrals of motion, namely

$$X_1 = P_1^2 - P_2^2 + \omega^2(x^2-y^2) + \frac{2b}{x^2} - \frac{2c}{y^2} \tag{3.7}$$

$$X_2 = L_3^2 + 2r^2(\frac{b}{x^2} + \frac{c}{y^2}).$$

The Hamiltonian H allows separation of variables in cartesian and polar coordinates and also in elliptic ones. We shall identify $X_1 = A$ and the cubic integral B is equal to

$$B = -\frac{1}{8}\{X_1, X_2\} = xP_1P_2^2 - yP_1^2P_2 + xy(\frac{-2b}{x^3} + \omega^2 x)P_2 - xy(\frac{-2c}{y^3} + \omega^2 y)P_1 \tag{3.8}$$

$$\{A,B\} = C = 4(A^2 - 2HA - H^2 + 8\omega^2 X_2 - 16\omega^2(b+c)$$
$$\{A,C\} = -64\omega^2 B \tag{3.9}$$
$$\{B,C\} = -2A^3 + 8H^2A + 64(c-b)\omega^2 H + 32(c+b)\omega^2 A$$
$$K = C^2 + 64\omega^2 B^2 - A^4 + (8H^2 + 32(c+b)\omega^2)A^2 + 128(c-b)\omega^2 HA \tag{3.10}$$

$$= 16H^4 - 128\omega^2(c+b)H^2 + 1024\omega^4 bc$$

In this case there is no underlying finite-dimensional Lie algebra, however X_1, X_2 and H generate a quadratic Lie algebra[16,18]

$$\begin{aligned} \{X_1, X_2\} &= C_q \\ \{X_1, C_q\} &= -8X_1^2 + 16X_1H - 16\omega^2 X_2 + 32\omega^2(b+c) \\ \{X_2, C_q\} &= 16X_1X_2 - 16HX_2 + 32H(c-b) \end{aligned} \tag{3.11}$$

With $C_q = -B$ and the relations (3.9) of the cubic algebra are consequences of (3.11)

Case 3. A further quadratically superintegrable Hamiltonian[5,6] is

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + \frac{\omega^2}{2}(4x^2 + y^2) + \frac{b}{y^2} + cx \tag{3.12}$$

The two second order integrals of motion are

$$A = P_1^2 - P_2^2 + \omega^2(4x^2 - y^2) - \frac{2b}{y^2} + 2cx \tag{3.13}$$

$$X_2 = xP_2^2 - yP_1P_2 + \frac{2bx}{y^2} - 2axy^2 - \frac{c}{2}y^2$$

and the Hamiltonian allows the separation of variables in cartesian and parabolic coordinates. The third order integral is

$$B = -\frac{1}{4}\{X_1, X_2\} = P_1P_2^2 + (\frac{2b}{y^2} - \omega^2 y^2)P_1 + (4\omega^2 xy + cy)P_2 \tag{3.14}$$

$$\begin{aligned} \{A, B\} &= C = 16\omega^2 X_2 \\ \{A, C\} &= -64\omega^2 B \\ \{B, C\} &= 16\omega^2 HA + 16\omega^2 H^2 - 12\omega^2 A^2 + 8c^2 H - 4c^2 A + 128\omega^4 b \end{aligned} \tag{3.15}$$

$$K = C^2+64\omega^2B^2-8\omega^2A^3+(16\omega^2H-4c^2)A^2+(32\omega^2H^2+16c^2H+256\omega^4b)A \tag{3.16}$$
$$= 64\omega^2H^3+16c^2H^2-512\omega^4bH-128\omega^2bc^2$$

As in Case 2 we have a cubic algebra but all relations in it follow from the already known quadratic algebra[16,18].

Case 4. We have

$$H = \frac{P_1^2}{2}+\frac{P_2^2}{2}+\frac{\omega^2}{2}(9x^2+y^2) \tag{3.17}$$

Which is anisotropic harmonic oscillator with a rational frequency ratio ω_1/ω_2=3/1. The second and third order integrals are

$$A = P_x^2-P_y^2+\omega^2(9x^2-y^2) \tag{3.18}$$

$$B = -yP_1P_2^2+xP_2^3+\frac{1}{3}\omega^2y^3P_1-3\omega^2xy^2P_2$$

$$\{A,B\} = C = -4P_1P_2^3+12\omega^2y^2P_1P_2-36\omega^2xyP_2^2+12\omega^4xy^3)$$
$$\{A,C\} = -144\omega^2B \tag{3.19}$$
$$\{B,C\} = 2A^3+8H^3-6HA^2$$
$$K = C^2+144\omega^2B^2+A^4-4HA^3+16AH^3 = 16H^4 \tag{3.20}$$

In this case the integral B is irreducible, i.e. it is not the Poisson bracket of lower order integrals. The cubic algebra (3.20) is related to a u(2) invariance algebra constructed by Jauch and Hill[1] for any anisotropic harmonic oscillator in E_2 with a rational frequency ratio. Their Lie algebra for the ratio ω_1/ω_2=3/1 is generated by the integrals.

$$\{H, \quad B_1=\frac{C}{H-A}, \quad B_2=\frac{B}{H-A}, \quad B_3=A\} \tag{3.21}$$

It has actually been shown that in many cases in classical mechanics functions of integrals of motion can be constructed that generate finite dimensional Lie algebras[25].In the case of the anisotropic oscillator (3.17) it suffices to take the fractions B_1 and B_2 of the polynomials (3.17),...,(3.20). The Hamiltonians considered so far namely (3.1),(3.6),(3.12) and (3.17) are all superintegrable, both in the classical and quantum cases[23].

The remaining 4 cases are quite different in that the systems are integrable only in the classical case. They are all obtained as singular limits of quantum integrable systems. As we shall see, the quantum systems are quite different and the potentials are expressed in terms of Painlevé transcendents.

Case 5. The classical Hamiltonian and two integrals of motion in this case are

$$H=\frac{P_1^2}{2}+\frac{P_2^2}{2}+\beta_1^2\sqrt{|x|}+\beta_2^2\sqrt{|y|}, A=\frac{P_1^2}{2}-\frac{P_2^2}{2}+\beta_1^2\sqrt{|x|}-\beta_2^2\sqrt{|y|} \quad (3.22)$$

$$B=\beta_2^4P_1^3+\epsilon\beta_1^4P_2^3+3\beta_2^4\beta_1^2\sqrt{|x|}P_1+\epsilon 3\beta_1^4\beta_2^2\sqrt{|y|}P_2$$

$$\begin{aligned}\epsilon&=1, xy>0 \\ \epsilon&=-1, xy<0 \quad .\end{aligned} \quad (3.23)$$

In the quantum case an integral of the type B exists if the potential V(x,y)=$V_1(x)+V_2(y)$ satisfies[23]

$$\begin{aligned}\hbar^2V_1''(x)&=6V_1^2(x)-6\beta_1^4x \\ \hbar^2V_2''(y)&=6V_2^2(y)-6\beta_2^4y\end{aligned} \quad (3.24)$$

The classical (and singular) limit $\hbar\to 0$ yields the potential in (3.22). The cubic algebra (2.5) in this case simplifies and we get

$$\begin{aligned}\{A,B\}&=C=3\beta_1^4\beta_2^4 \\ \{A,C\}&=0 \\ \{B,C\}&=0\end{aligned} \quad (3.25)$$

$$K = 9\beta_1^8\beta_2^8 \tag{3.26}$$

Thus $\{A, B, C\}$ actually generate a nilpotent Lie algebra, isomorphic to the Heisenberg algebra in one dimension (since C is constant).

We can see from eq. (3.24) that the quantum case will be completely different. Indeed eq. (3.24) can be solved in terms of the first Painlevé transcendent ${P_I}^{24}$.

Case 6. The classical Hamiltonian and two integrals of motion are

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + \frac{\omega^2}{2}y^2 + V(x), A = \frac{P_1^2}{2} - \frac{P_2^2}{2} - \frac{\omega^2}{2}y^2 + V(x) \tag{3.27}$$

$$B = -yP_1^3 + xP_1^2P_2 + (\frac{\omega^2}{2}x^2 - 3V)yP_1 - \frac{1}{\omega^2}(\frac{\omega^2}{2}x^2 - 3V)V_xP_2$$

where V satisfies a quartic equation

$$-9V(x)^4 + 14\omega^2x^2V(x)^3 + (6d - 3\frac{\omega^4}{4}x^4)V(x)^2 + (\frac{3\omega^6}{2}x^6 - 2\omega^2x^2)V(x) \tag{3.28}$$

$$+(cx^2 - d - d\frac{\omega^4}{2}x^4 - \frac{\omega^8}{16}x^8) = 0$$

In the quantum case V satifies a fourth order differential equation[23]

$$\hbar^2V^{(4)} = 12\omega^2xV' + 6(V^2)'' - 2\omega^2x^2V'' + 2\omega^4x^2 \tag{3.29}$$

that can be solved in terms of the Painlevé transcendent ${P_{IV}}^{24}$. Eq. (3.27) is the solution of (3.29) for $h \rightarrow 0$ and c and d are integration constants. In general, eq. (3.27) has 4 roots and the expressions for them are quite complicated. A special case occurs if ω^2,c and d satisfy.

$$c = \frac{2^3\omega^8b^3}{3^6}, d = \frac{\omega^4b^2}{3^3} \tag{3.30}$$

where b is an arbitrary constant. Then eq. (3.28) has a double root and we obtain

$$V_{1,2}(x) = \frac{\omega^2}{18}(2b + 5x^2 \pm 4x\sqrt{b + x^2}) \tag{3.31}$$

$$V_3(x) = V_4(x) = (-\frac{\omega^2 b}{3^3} + \frac{\omega^2}{2}x^2) \tag{3.32}$$

For V(x) satisfying (3.28) the cubic algebra is

$$\begin{aligned} \{A,B\} &= C \\ \{A,C\} &= -4\omega^2 B \\ \{B,C\} &= 8A^3 + 12HA^2 - 4H^3 - 4\frac{4b^2\omega^4}{27}A + \frac{4b^3\omega^6}{729} \end{aligned} \tag{3.33}$$

$$\begin{aligned} K &= C^2 + \omega^2 B^2 + 4A^4 + 8HA^3 - 4bA^2 + (-8H^3 + \frac{8b^3\omega^6}{729})A \\ &= 4H^4 - \frac{4}{27}b^2\omega^4 H^2 + \frac{8b^3\omega^6}{729}H \end{aligned} \tag{3.34}$$

Case 7. The classical Hamiltonian and two integrals of motion in this case are

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + a^2|y| + b^2\sqrt{|x|}, A = \frac{P_1^2}{2} - \frac{P_2^2}{2} - a^2|y| + b^2\sqrt{|x|} \tag{3.35}$$

$$B = P_1^3 + 3b^2\sqrt{|x|}P_1 + \epsilon\frac{3b^4}{2a^2}P_2$$

$$\begin{aligned} \epsilon &= -1, xy > 0 \\ \epsilon &= 1, xy < 0 \quad . \end{aligned} \tag{3.36}$$

In the quantum case the potential is V(x,y)=ay + V(x) where V(x) satisfies

$$\hbar^2 V'' = 6V^2 - 6b^4 x, \qquad b \neq 0 \tag{3.37}$$

This is solved in terms of the first Painlevé transcendent for $\hbar \neq 0$. For $\hbar = 0$ the singular limit is V=$b\sqrt{x}$. The cubic algebra in this case is very simple and reduces to a Lie algebra, the Heisenberg algebra as in Case 5. We have

$$\{A,B\} = C = 3b^4$$

$$
\begin{aligned}
\{A, C\} &= 0 && (3.38)\\
\{B, C\} &= 0\\
K &= 9b^8 && (3.39)
\end{aligned}
$$

Case 8. In this case we have

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + ay + V(x), \quad A = \frac{P_1^2}{2} - \frac{P_2^2}{2} - ay + V(x) \tag{3.40}$$

$$B = aP_1^3 - bP_1^2P_2 + a(3V(x) - bx)P_1 - 2bV(x)P_2$$

where V(x) satisfies a cubic equation

$$V(x)^3 - 2bxV(x)^2 + b^2x^4V(x) - d = 0 \tag{3.41}$$

This is again obtained from a singular limit of a second order nonlinear ODE in the quantum case. It is, for $\hbar \neq 0$, solved in terms of the Painlevé transcendent P_{II}. The cubic algebra in this case reduces to

$$
\begin{aligned}
\{A, B\} &= C = 2ab(A + H)\\
\{A, C\} &= 0 && (3.42)\\
\{B, C\} &= -4a^2b^2(A + H)\\
K &= 4a^2b^2H^2 && (3.43)
\end{aligned}
$$

We see that (3.42) is actually a Lie algebra with centre H. If we put

$$A_1 = \frac{H + A}{2}, B_1 = -\frac{1}{2ab}B \tag{3.44}$$

We obtain the solvable decomposable Lie algebra

$$\{H_1, B_1\} = B_1, \{H_1, H\} = \{B_1, H\} = 0 \tag{3.45}$$

The cubic equation (3.41) can be taken to standard form by putting

$$V(x) = y(x) + \frac{2bx}{3} \tag{3.46}$$

We obtain

$$y^3 + 3py + 2q = 0, \quad p = -(\frac{bx}{3})^2, \quad q = (\frac{bx}{3})^3 - \frac{d}{2} \tag{3.47}$$

This discriminant is

$$D = q^2 + p^3 = \frac{d^2}{4} - \frac{b^3 dx^3}{27} \tag{3.48}$$

For D $<$ 0,i.e. $x^3 > \frac{27d}{4b^2}$ with d$<$0 or $x^3 < \frac{27d}{4b^2}$ with d$>$0 it has 3 real roots,
For D $>$ 0,i.e. $x^3 < \frac{27d}{4b^2}$ with d$<$0 or $x^3 > \frac{27d}{4b^2}$ with d$>$0 it has 1 real root y_1 and 2 complex ones,$y_3 = \bar{y}_2$.

The corresponding potentials are

$$\begin{aligned} V_1(x) = \frac{2bx}{3} &+ \frac{2^{1/3} b^2 x^2}{3(2 + d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}} \\ &+ \frac{(27d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}}{32^{1/3}} \end{aligned} \tag{3.49}$$

$$\begin{aligned} V_2(x) = \frac{2bx}{3} &+ \frac{(1 + i\sqrt{3}) b^2 x^2}{32^{2/3}(2 + d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}} \\ &+ \frac{(1 - i\sqrt{3})(27d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}}{62^{1/3}} \end{aligned} \tag{3.50}$$

$$\begin{aligned} V_3(x) = \frac{2bx}{3} &+ \frac{(1 - i\sqrt{3}) b^2 x^2}{32^{2/3}(2 + d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}} \\ &+ \frac{(1 + i\sqrt{3})(27d - 2b^3x^3 + 3\sqrt{3}\sqrt{27d^2 - 4b^3dx^3})^{1/3}}{62^{1/3}} \end{aligned} \tag{3.51}$$

Multiple roots occur for D=0.This happens (for x $\neq$ const) for d=0 only and then we have

$$V = bx \tag{3.52}$$

as a double root(and V=0 as the simple one). We can consider the particular Hamiltonian,

$$H = \frac{P_1^2}{2} + \frac{P_2^2}{2} + a^2|y| + b^2|x|, A = \frac{P_1^2}{2} - \frac{P_2^2}{2} - a^2|y| + b^2|x| \tag{3.53}$$

$$B = a^2P_1^3 - b^2P_1^2P_2 + 2a^2b^2|x|P_1 - 2b^4|x|P_2$$

4 Trajectories for classical superintegrable systems

If a Hamiltonian system is maximally superintegrable and satisfies certain analyticity properties, then all of its bounded trajectories are closed and the motion is periodic[26]. Here we shall discuss the trajectories for all the 8 systems of Section 3. The trajectories can be obtained in a uniform manner directly from the integrals of motion. Indeed in all cases we can put

$$\frac{1}{2}P_1^2 + f(x) = E_1 \tag{4.1}$$

$$\frac{1}{2}P_2^2 + g(y) = E_2 \tag{4.2}$$

$$B = \mu P_1^3 + \nu P_1^2 P_2 + \rho P_1 P_2^2 + \sigma P_2^3 + \phi P_1 + \psi P_2 = k \quad . \tag{4.3}$$

In (4.3) μ ,ν ,ρ and σ are low order polynomials in x and y, ϕ and ψ are functions of x and y (all of them known). The constants E_1 and E_2 are positive, k arbitrary. From (4.1) and (4.2) we obtain P_1^2 and P_2^2 in term of x and y, respectively. From eq(4.3) we obtain

$$\begin{aligned}&(P_1^2(\mu P_1^2 + \rho P_2^2 + \phi)^2 - P_2^2(\nu P_1^2 + \sigma P_2^2 + \psi)^2)^2 + k^4 \qquad (4.4)\\ &= 2k^2(P_1^2(\mu P_1^2 + \rho P_2^2 + \phi)^2 + P_2^2(\nu P_1^2 + \sigma P_2^2 + \psi)^2) \quad .\end{aligned}$$

Substituting for P_1^2 and P_2^2 from (4.1) and (4.2) we obtain the equation for the trajectories. Directly from eq (4.1) and (4.2) we see that the motion is bounded if there exist two constants, x_o and y_o,such that we have

$$f(x) \geq 0, \quad x^2 > x_o^2, \quad g(y) \geq 0, y^2 > y_o \quad . \tag{4.5}$$

This direct method of computing trajectories is universal, hence not necessarily convenient in any special case. For instance for the harmonic oscillator it gives a polynomial of order 16 in x and y which can then be simplified to a second order one. However it avoids the problem of integrating the equation of motion and then eliminating time.

Case 1. The trajectories of the harmonic oscillator are well known and are ellipses.

Case 2. The trajectories are given in[6] and are easily obtained by integrating the equation of motion :

$$x(t)^2 = \frac{E_1}{\omega^2} + \sqrt{\frac{E_1^2}{2\omega^4} - \frac{2b}{\omega^2}} sin(2\omega t + c_1) \tag{4.6}$$

$$y(t)^2 = \frac{E_2}{\omega^2} + \sqrt{\frac{E_2^2}{2\omega^4} - \frac{2c}{\omega^2}} sin(2\omega t + c_2) \tag{4.7}$$

Case 3. The trajectories are also given in[6] and are obtained as in Case 2.

$$x(t)^2 = \frac{-c}{4\omega^2} + \sqrt{\frac{E_1}{\omega^2} + \frac{c^2}{32\omega^2}} sin(2\omega t + c_1) \tag{4.8}$$

$$y(t)^2 = \frac{E_2}{\omega^2} + \sqrt{\frac{E_2^2}{\omega^4} - \frac{2c}{\omega^2}} sin(2\omega t + c_2) \tag{4.9}$$

Case 4.The Case of the anisotropic oscillator is well known. The trajectories are what is called Lissajous figures.

$$x(t) = \sqrt{\frac{2E_1}{9\omega^2}} sin(3\omega t + c_1) \tag{4.10}$$

$$y(t) = \sqrt{\frac{2E_2}{\omega^2}} sin(\omega t + c_2) \tag{4.11}$$

Case 5. The potential is

$$V = \beta_1^2 \sqrt{|x|} + \beta_2^2 \sqrt{|y|} \quad . \tag{4.12}$$

We find the trajectories using the equation (4.4). The bounded trajectories are closed. Examples are shown on Fig 5.

Case 6. We know from the Section 3 that the only particular case where two roots coincide is (3.30). The trajectories for the potentials $V_{1,2}(x)$ satisfy

$$y(t) = \sqrt{\frac{2E_2}{\omega^2}} sin(\omega t + c_2) \tag{4.13}$$

$$\int \frac{dx}{\sqrt{-\frac{10\omega^2}{9}x^2 \mp \frac{4\omega^2}{9}x\sqrt{b+x^2} + 2E_1}} = t + c_1 \tag{4.14}$$

The trajectories were obtained numerically directly from the equations of motion. We have closed trajectories and examples are shown on Fig 6. For $V_3(x)$ the potential is a shifted harmonic oscillator, so the trajectories are ellipses.

Case 7. The potential is

$$V = a^2|y| + b^2\sqrt{|x|} \tag{4.15}$$

We find the trajectories using equation (4.4). The bounded trajectories are closed. They are shown on Fig 7.

Case 8. There is the particular case

$$V = a^2|y| + b^2|x| \tag{4.16}$$

We use eq. (4.4) to calculate the trajectories. Again the bounded trajectories are closed. An example is shown on Fig 8.

5 Conclusion

The main results of this article are the study of the algebras of integrals of motion, presented in Sections 2 and 3 and the investigation of classical trajectories in Section 4.

In all 8 cases of superintegrable systems, separating in cartesian coordinates and allowing a third order integral of motion, the integrals of motion generate a cubic Poisson algebra. In many cases this polynomial algebra is reducible that is it is a consequence of the existence of a simpler algebraic structure.

In 4 cases the simplest underlying structure is a Lie algebra. Thus in Case 1, the isotropic harmonic oscillator, the integrals of motion generate the u(2) algebra with the Hamiltonian as its central element. The third order operator B and the fourth order one C lie in the enveloping algebra of u(2).

In Case 5 the potential $V(x,y) = \beta_1^2\sqrt{|x|} + \beta_2^2\sqrt{|y|}$ is nonanalytical at the origin. In this case the Poisson bracket of the second and the third order integrals is a constant. Hence A,B and C generate the Heisenberg Lie algebra.

Case 7 with $V(x,y) = a^2|y| + b^2\sqrt{|x|}$ is similar to Case 5 in that A, B and C generate the Heisenberg Lie algebra (with C constant). In both of these cases the Lie algebra is actually

$$\{H\}\bigoplus\{A,B,C\} \tag{5.1}$$

i.e. the direct sum of a one-dimensional Lie algebra, generated by the Hamiltonian, and the Heisenberg algebra.

In Case 8 we have V(x,y)= ay + V(x) with V(x) satisfying the cubic equation (3.41), the integrals generate a solvable decomposable Lie algebra (3.45). The Hamiltonian H commutes with the elements of the two-dimensional solvable algebra.

The Case 2 and 3 are different. The Hamiltonians (3.6) and (3.12) are actually quadratically superintegrable and the third order integral B is the Poisson bracket of two second order ones. These second order integrals give rise to a quadratic Poisson algebra[15,16] and the cubic algebra of our Section 3 is an algebraic consequence of the quadratic one.

Finally, two irreducible cases remain. One is the Case 4, the anisotropic harmonic oscillator (3.17). We have a genuine third order polynomial algebra (3.19), (3.20) of polynomial integrals H,A,B and C of order 2,2,3 and 4, respectively. Somewhat artificially, we can construct the Lie algebra u(2) of eq. (3.21), using rational functions of these integrals.

Case 6 is again irreducible, i.e. we obtain the genuinely cubic polynomial algebra (3.33), (3.34), again involving two second order, one third order and one fourth order integral. The condition for this algebra to close is precisely that V(x) should satisfy the quartic equation (3.28).

To our knowledge, no classification of polynomial algebras exist, not even of quadratic ones, still less of cubic algebras. We can however see that the linear parts of the algebras of Case 4 and Case 6 are not isomorphic as Lie algebras. That of Case 4 is solvable, that of Case 6 is simple.

Trajectories are studied in Section 4 with examples on Fig 1,...,8. The most important result is that all finite trajectories are closed. We note that the potentials of Case 1 and 4 are analytic everywhere, so superintegrability implies periodic motion[26]. In cases 2 and 3 the potentials are singular along one or both coordinate axes, analytic elsewhere. In Case 5 the potential is nonanalytic with nonisolated branchpoints along the axes. That not withstanding, bounded trajectories turn out to be closed.

The functions $V_{1,2}(x)$ in Case 6 are nonanalytic for $x^2 = -b$ for b>0. Similar statements apply for Case 7. In Case 8 we have studied a psecial case when the cubic equation (3.41) has a double root. In this case we either have a linear potential $V(x,y) = \alpha x + \beta y$ with no bounded trajectories, or the nonanalytical one $V(x,y) = a^2|y| + b^2|x|$ (4.16) that we have investigated. Our conclusion is that the bounded trajectories are always closed for the superintegrable potentials, wheter they are analytical, or not.

An investigation of the properties of superintegrable systems with a third order integral of motion is in progress.

Acknowledgments The authors thanks Frederick Tremblay for useful discussions. I.M. benefited from an FQRNT student fellowship.

6 References

[1]J.M Jauch and E.L Hill,On the problem of degeneracy in quantum mechanics,Phys Rev,57,641-645 (1940).

[2] M.Moshinsky and Yu F.Smirnov,The Harmonic Oscillator In Modern Physics,Harwood,Amsterdam,641-645 (1966).

[3] V. Fock, Zur Theorie des Wasserstoffsatoms Phys.98,145-154 (1935).

[4] V. Bargmann,Z. Zur Theorie des Wasserstoffsatoms Phys. 99,576-582 (1936).

[5] J. Fris, V. Mandrosov, Ya.A. Smorodinsky, M. Uhlir and P. Winternitz, On higher symmetries in quantum mechanics, Phys. Lett. 16, 354-356 (1965).

[6] P.Winternitz, Ya.A Smorodinsky, M.Uhlir and I.Fris, Symmetry group in classical and quantum mechanics , J.Nucl.Phys.(U.SS.R)4,625-635 (1966).

[7] A. Makarov, Ya.A. Smorodinsky, Kh. Valiev and P. Winternitz,A systematic search for nonrelativistic systems with dynamical symmetries, Nuovo Cim. A52, 1061-1084 (1967).

[8] E.G. Kalnins,J.M.Kress and W.Miller Jr, Second-order superintegrable systems in conformally flat spaces. I,II,III, J.Math.Phys,46,053509 28 pages,053510 15 pages,103507 28 pages,(2005).

[9] E.G. Kalnins, W. Miller Jr. and G. S. Pogosyan. Completeness of multiseparability in E2,C. J. Phys. A33,4105-4120 (2000).

[10] M.B.Sheftel, P.Tempesta and P.Winternitz,Exact solvability of superintegrable systems, J.Math.Phys.42,659-673 (2001).

[11] P.Tempesta, A.V Turbiner and P.Winternitz,Exact solvability of superintegrable systems, J.Math.Phys.42,4248-4257 (2001).

[12] N.W. Evans. Superintegrability in classical mechanics. Phys. Rev. A 41,5666-5676 (1990).

[13] N.W. Evans. Group theory of the Smorodinsky-Winternitz system. J. Math. Phys. 32,3369-3375 (1991).

[14] Ya. I Granovskii,A.S. Zhedanov and I.M. Lutzenko , Quadratic Algebra

as a Hidden Symmetry of the Hartmann Potential, J. Phys. A24 3887-3894 (1991).

[15]P.Létourneau and L.Vinet, Superintegrable systems :Polynomial algebras and quasi-exactly solvable hamiltonians,Ann. Physics 243,144-168(1995).

[16]C.Daskaloyannis,Quadratic poisson algebras of two-dimensional classical superintegrable systems and quadratic associative algebras of quantum superintegrable systems, J.Math.Phys.42,1100-1119, (2001).

[17]D.Bonatsos,C.Daskaloyannis and K.Kokkotas, Deformed oscillator algebras for two-dimensional quantum superintegrable systems. Phys. Rev.A 50,3700-3709 (1994).

[18]D.Bonatsos,C.Daskaloyannis and K.Kokkotas, Quantum-algebraic description of quantum superintegrable systems in two dimensions. Phys. Rev. A 48,R3407-R3410. 81S05 (1993).

[19]J. Drach Sur l'intégration logique des équations de la dynamique à deux variables :Forces conservatrices.Intégrales cubiques.Mouvements dans le plan,C.R. acad. Sci III,200,22-26 (1935).

[20]J. Drach, Sur l'intégration logique et sur la transformation des équations de la dynamique à deux variables :Forces conservatrices.Intégrales.C.R.Acad.Sci III,599-602 (1935).

[21]J.Hietarinta,Classical vs. Quantum integrability,J.Math.Phys.25,1833-1840 (1984).

[22]S.Gravel and P.Winternitz, Superintegrability with third-order integrals in quantum and classical mechanics J. Math. Phys. 43 5902-5912 (2002).

[23]S.Gravel, Hamiltonians separable in Cartesian coordinates and thirdorder integrals of motion, J. Math. Phys. 45 1003-1019 (2004).

[24]E.L.Ince,Ordinary Differential Equations,Dover,New York, (1956).

[25]G. Gonora,P. Kosinski,M. Majenski and P.Mashenka,Symmetry algebras

for superintegrable systems.J.Phys.A,39,343-349 (2006).

[26]N.N.Nekhoroshev,Action-angle variables and their generalizations.Trudy Mskov.Mat.Obshch.26,181-198 (1972).

Figure captions

Fig1.A trajectory for $V = \frac{\omega^2}{2}(x^2 + y^2)$. Parameter $\omega^2 = 2$, $v_{xo} = -1.5$, $x_o = 5$, $v_{yo} = -1.2$, $y_o = -2$, t=[0,400]

Fig2.A trajectory for $\frac{\omega^2}{2}(x^2 + y^2) + \frac{b}{x^2} + \frac{c}{y^2}$. Parameter $\omega^2 = 2$,b=2 and c=3, $v_{xo} = -1.5$, $x_o = 5$, $v_{yo} = -1.2$, $y_o = -2$, t=[0,400]

Fig3.A trajectory for $\frac{\omega^2}{2}(4x^2 + y^2) + \frac{b}{y^2} + cx$. Parameter $\omega^2 = 1$,b=2 and c=3, $v_{xo} = -1.5$, $x_o = 5$, $v_{yo} = -1.2$, $y_o = -2$, t=[0,400]

Fig4.A trajectory for $\frac{\omega^2}{2}(9x^2 + y^2)$. Parameter $\omega^2 = 1$ and d=3, $v_{xo} = -1.5$, $x_o = 5$, $v_{yo} = -1.2$, $y_o = -2$, t=[0,400]

Fig5.A trajectory for $V = \beta_1^2\sqrt{|x|} + \beta_2^2\sqrt{|y|}$. Parameter $E_1 = E_2 = k = 1$

Fig6.A trajectory for $\frac{\omega^2}{18}(2b + 5x^2 \pm 4x\sqrt{b + x^2})$. Parameter $\omega^2 = 1$ and d=3, $v_{xo} = -1.5$, $x_o = 5$, $v_{yo} = -1.2$, $y_o = -2$, t=[0,400]

Fig7.A trajectory for $V = a^2|y| + b^2\sqrt{|x|}$. Parameter $E_1 = E_2 = k = 1$

Fig8.A trajectory for $V = a^2|y| + b^2|x|$. Parameter $E_1 = E_2 = k = 1$


2.5
2
1.5
1
0.5
0
−0.5
−1
−1.5
−2
−2.5
−6
−4
−2
0
2
4
6


-0.6
-0.8
-1
-1.2
-1.4
-1.6
-1.8
-2
-2.2
-2.4
-2.6
0
1
2
3
4
5
6


−0.6
−0.8
−1
−1.2
−1.4
−1.6
−1.8
−2
−2.2
−2.4
−2.6
−6
−4
−2
0
2
4
6


2.5
2
1.5
1
0.5
0
−0.5
−1
−1.5
−2
−2.5
−6
−4
−2
0
2
4
6


1
0.5
0
-0.5
-1
-1
-0.5
0
0.5
1



2.5
2
1.5
1
0.5
0
−0.5
−1
−1.5
−2
−2.5
−20
−15
−10
−5
0
5
10


1
0.5
0
-0.5
-1
-1
-0.5
0
0.5
1


1
0.5
0
-0.5
-1
-1
-0.5
0
0.5
1